

## Tilburg University

### **An asymptotic justification for a modified GLS procedure to estimate ARMA parameters**

van der Leeuw, J.L.; Tigelaar, H.H.

*Publication date:*  
1994

[Link to publication in Tilburg University Research Portal](#)

*Citation for published version (APA):*

van der Leeuw, J. L., & Tigelaar, H. H. (1994). *An asymptotic justification for a modified GLS procedure to estimate ARMA parameters*. (Research memorandum / Tilburg University, Department of Economics; Vol. FEW 662). Unknown Publisher.

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CBM

762R

7626

1994

NR.662




## Research Memorandum

Faculty of Economics and  
Business Administration

Tilburg University





**AN ASYMPTOTIC JUSTIFICATION  
FOR A MODIFIED GLS  
PROCEDURE TO ESTIMATE  
ARMA PARAMETERS**

Jan van der Leeuw  
Harry Tigelaar

**FEW 662**

Communicated by Prof.dr. B.B. van der Genugten

AN ASYMPTOTIC JUSTIFICATION FOR A MODIFIED GLS PROCEDURE  
TO ESTIMATE ARMA PARAMETERS

Jan van der Leeuw and Harry Tigelaar  
Dept. of Econometrics  
Tilburg University  
P.O.Box 90153  
NL - 5000 LE Tilburg

**Abstract**

The asymptotic form of the concentrated likelihood function of a stationary ARMA model is investigated. Basis for the analysis is the closed form formula for the exact covariance matrix. GLS estimates of regression coefficients can be based on a simple and widely used adjusted form of the covariance matrix. However, ML-estimates of the ARMA parameters become more complicated, as they depend on the limit of the determinant of the covariance matrix.

JEL code: C22

Keywords : Generalised Linear Model; Autoregressive moving average process; exact ARMA covariance matrix.

## 1. Introduction

Several authors have presented procedures to estimate the covariance parameters, be it for a pure time series model (Kohn and Ansley, 1985) or for a regression model (Zinde-Walsh and Galbraith, 1991). The resulting formulas and algorithms become very complicated. In Van der Leeuw (1993) relatively simple expressions are presented, based on a closed form of the general ARMA covariance matrix. This is possible as the closed form of the covariance matrix is simple enough to be differentiated analytically. The price to be paid for this procedure is the computation of the covariance matrix and its inverse. Moreover, second derivatives need very much (computer) time consuming algorithms, as they depend on the complete covariance matrix.

In this paper the asymptotic form of the covariance matrix and its determinant are studied. Maximum likelihood estimation of the ARMA model is equivalent to determining the first order condition of the concentrated likelihood. In this expression the covariance matrix appears twice: as determinant and as part of a quadratic form. We will show what happens with both forms when the number of observations becomes large.

## 2. The linear model

Consider the linear model  $y = X\beta + \varepsilon$ , where  $y$  has dimensions  $(T \times 1)$ ,  $X$   $(T \times k)$ ,  $\beta$   $(k \times 1)$  and  $\varepsilon$   $(T \times 1)$  and covariance matrix  $E\varepsilon\varepsilon^T = \sigma^2 V$ , which is a function of a vector  $\vartheta$ . One way to estimate the unknown parameters  $\beta$  and  $\vartheta$  is by minimizing the weighted sum of squares  $\varepsilon^T V^{-1} \varepsilon$ . Supposing normally distributed errors, we may prefer to maximize the (concentrated) likelihood function, which is equivalent to minimizing  $S = |V|^{1/T} e^T V^{-1} e$  (Judge *et al.*, p.284). Here  $e = y - Xb$ , where  $b$  is the Aitken estimator of  $\beta$ .

Minimizing  $S=|V|^{1/T} e^T V^{-1} e$  is equivalent to solving the first order conditions or to find a solution to

$$dS=|V|^{1/T} \{s^2 \text{tr} V^{-1} dV + e^T d(V^{-1})e\} = 0 \quad (1)$$

with  $s^2 = e^T V^{-1} e / T$ . In the next section we show how  $V$  can be written in a form that can be differentiated easily and substituted into (1). The pure time series model is similar:  $e$  need not to be computed as it is identical to  $y$ .

### 3. The ARMA covariance matrix

The general form of ARMA distributed errors is given by

$$\varepsilon_t = - \sum_{i=1}^p \vartheta_i \varepsilon_{t-i} + v_t + \sum_{i=1}^q \alpha_i v_{t-i}, \quad t=1, \dots, T, \quad (2)$$

where  $v_t$  is a sequence of independently and identically distributed random variables.  $\vartheta$  denotes the vector  $(\vartheta_1, \vartheta_2, \dots, \vartheta_p)^T$  of AR-parameters,  $\alpha$  is the vector  $(\alpha_1, \alpha_2, \dots, \alpha_q)^T$  of MA-parameters. We assume that  $p$  and  $q$  are known and that the invertibility conditions are fulfilled. By definition  $\vartheta_0$  and  $\alpha_0$  are put equal to 1. Use  $\sigma^2 V$  to denote the covariance matrix of  $\varepsilon$ :

$$\sigma^2 V = E \varepsilon \varepsilon^T.$$

Following Pagan (1974), we introduce two matrices for both the AR parameters and the MA parameters. We define a (square) lower band matrix  $P$  of dimensions  $T \times T$ , and a  $T \times p$  matrix  $Q$  as follows:

$$P = \begin{bmatrix} 1 & & & & \\ \vartheta_1 & & & & \\ & \ddots & & & \\ & & \ddots & & \\ \vartheta_p & & & & \\ & \ddots & & & \\ & & \vartheta_p & & \\ & & & \ddots & \\ & & & & \vartheta_1 & 1 \end{bmatrix}, \quad Q = \begin{bmatrix} \vartheta_p & \vartheta_{p-1} & \dots & \vartheta_1 \\ 0 & & & \\ & \ddots & & \\ 0 & & & \vartheta_p \\ 0 & & & 0 \\ & \ddots & & \\ 0 & & & 0 \end{bmatrix}.$$

The upper triangular part of a lower band matrix consists of zeros and the lower part has off-diagonals with the same elements.  $Q$  consists of an upper



$p \times p$  part with an upper band matrix and a lower  $(T-p) \times p$  part, which consists of only zeros. Like  $P$  and  $Q$  will be used to describe the AR part of the error vector, so are  $M$  and  $N$  defined for the MA part, where  $\theta$  is replaced by  $\alpha$  and  $p$  by  $q$ .

Defining the auxiliary vectors  $\bar{\varepsilon} = (\varepsilon_{-p+1}, \varepsilon_{-p+2}, \dots, \varepsilon_{-1}, \varepsilon_0)^T$  and  $\bar{v} = (v_{-q+1}, v_{-q+2}, \dots, v_{-1}, v_0)^T$ , we can write (2) in matrix form:

$$\begin{bmatrix} Q & P \end{bmatrix} \begin{bmatrix} \bar{\varepsilon} \\ \varepsilon \end{bmatrix} = \begin{bmatrix} N & M \end{bmatrix} \begin{bmatrix} \bar{v} \\ v \end{bmatrix}.$$

As is proven elsewhere (Van der Leeuw, 1992) in the exact covariance matrix,  $E\varepsilon\varepsilon^T = \sigma^2 V$ , of ARMA errors,  $V$  is equal to  $[N \ M] [\bar{P}^T \bar{P} - \bar{Q} \bar{Q}^T]^{-1} [N \ M]^T$ , where  $\bar{P}$  is like  $P$ , but of order  $(T+p) \times (T+p)$  and  $\bar{Q}$  like  $Q$ , but of order  $(T+p) \times p$ . In the MA case this expression reduces to  $[N \ M][N \ M]^T$  and in the AR case it becomes  $[P^T P - Q Q^T]^{-1}$ . When we use the underlined symbols  $\underline{P}$ ,  $\underline{Q}$ ,  $\underline{M}$ , and  $\underline{N}$ , we mean the upper left  $(p \times p)$  submatrix of corresponding matrix. Furthermore we will assume  $p$  greater than or equal to  $q$ . This gives no loss of generality as the case  $q \geq p$  is similar.

#### 4. Limits of quadratic form and determinant

As stated before the first order conditions for the AR and MA parameters are derived from the differential  $dS = |V|^{1/T} \{s^2 \text{tr} V^{-1} dV + e^T d(V^{-1}) e\}$ . Solving these conditions gives values for the parameters (see Van der Leeuw, 1993). However the computations will become very time consuming when the number of observations is large. This leads to the question whether a more efficient asymptotic form can be found. The asymptotic behaviour depends on the determinant part  $(\text{tr} V^{-1} dV)$  and the quadratic form  $e^T d(V^{-1}) e$ . Before we give some theorems regarding these limits we give a lemma about the inverse of the matrix  $M$  and a lemma for the asymptotic behaviour of a quadratic form to be used in the sequel.



### Lemma 1

The inverse of the matrix  $M$  can (apart of the last rows) be written in block form where the  $(i,j)^{th}$  block,

$$M_{i,j}^{-1} = \begin{cases} \underline{M}^{-1} & \text{if } i=j \\ (-\underline{M}^{-1}\underline{N})^{i-j}\underline{M}^{-1} & \text{if } i \geq j \\ 0 & \text{if } i < j \end{cases}$$

$\underline{M}$  refers to the  $p \times p$  upper left submatrix of  $M$ ,  $\underline{N}$  to the  $p \times p$  upper part of  $N$ . The eigenvalues of  $\underline{M}^{-1}\underline{N}$  are less than 1 in absolute value if (and only if) the invertibility condition holds.

### Proof

Observe that  $M$  can be written as  $\begin{bmatrix} \underline{M} & & \\ \underline{N} & \underline{M} & \\ & \underline{N} & \underline{M} \\ & & \ddots & \ddots \\ & & & \ddots & \ddots \end{bmatrix}$  where  $\underline{M}$  and  $\underline{N}$  are as defined

above. Direct verification gives the result. For the eigenvalues, see Van der Leeuw (1992).  $\square$

### Lemma 2

$$\lim_{T \rightarrow \infty} \text{tr} \{ (PM^{-1}N-Q)(P^T P - QQ^T)^{-1} (PM^{-1}N-Q)^T \} / T = 0.$$

### Proof

In the expression  $\left( \frac{PM^{-1}N-Q}{\sqrt{T}} \right) (P^T P - QQ^T)^{-1} \left( \frac{PM^{-1}N-Q}{\sqrt{T}} \right)^T$  the  $(p \times p)$  matrix  $(P^T P - QQ^T)$

depends on the AR-parameters and is independent of  $T$ . It is non singular (in fact positive definite) because of the invertibility condition.

Therefore we have to show that  $\lim_{T \rightarrow \infty} (PM^{-1}N-Q)/\sqrt{T} = 0$ .

The  $(T \times p)$  matrix  $PM^{-1}N-Q$  can be written as

$$(PM^{-1}N-Q) \begin{bmatrix} I_p \\ (-\underline{M}^{-1}\underline{N})^1 \\ (-\underline{M}^{-1}\underline{N})^2 \\ \vdots \\ (-\underline{M}^{-1}\underline{N})^i \\ \vdots \end{bmatrix}, \text{ as we will show now. Because } N \text{ is } \begin{bmatrix} \underline{N} \\ 0 \end{bmatrix}, M^{-1}N \text{ is a } (T \times p)$$

block column matrix of which the first element is  $\underline{M}^{-1}\underline{N}$ . The  $i^{\text{th}}$  block is

$$(-\underline{M}^{-1}\underline{N})^{i-1}\underline{M}^{-1}\underline{N}=(-1)^{i-1}(\underline{M}^{-1}\underline{N})^i, \quad i=2,3,\dots. \text{ Since } P=\begin{bmatrix} \underline{P} & \underline{P} \\ \underline{Q} & \underline{Q} \end{bmatrix}, \text{ the first}$$

$(p \times p)$  block of  $PM^{-1}N-Q$  becomes  $\underline{PM}^{-1}\underline{N}-\underline{Q}$ , and for subsequent blocks we get:

$$\underline{Q}(-1)^{i-2}(\underline{M}^{-1}\underline{N})^{i-1}+\underline{P}(-1)^{i-1}(\underline{M}^{-1}\underline{N})^i, \quad i=2, 3, \dots$$

or

$$(\underline{PM}^{-1}\underline{N}-\underline{Q})(-\underline{M}^{-1}\underline{N})^i, \quad i=1,2,\dots$$

From the invertibility condition we know, that  $|\lambda \underline{M}+\underline{N}|$  can only be zero if

$|\lambda|<1$ . This implies, that the eigenvalues of  $\underline{M}^{-1}\underline{N}$  are less than one in

absolute value and  $\lim_{T \rightarrow \infty} (\underline{M}^{-1}\underline{N})^T = 0$ . But this means, that every block of

$PM^{-1}N-Q$  times  $1/\sqrt{T}$  tends to zero.  $\square$

When  $T$  becomes large enough the influence of the 'starting' values of  $\bar{\epsilon}$  and  $\bar{v}$  will become less and less important. This implies that the role of the matrices  $N$  and  $Q$  can be neglected. This has been proven by several authors (e.g. Zinde-Walsh, 1992) in a more general context. Here we give a direct proof. It may be more clear if we write  $V$  as

$W+(N-MP^{-1}Q)(P^T P-QQ^T)^{-1}(N-MP^{-1}Q)^T$ , where  $W=P^{-1}M(P^{-1}M)^T$ . We want to show that asymptotically only  $W$  is important in estimating  $\beta$  and the ARMA parameters.

However, the determinant of  $V$  does not reduce to the determinant of  $W$  (which is one). This will be studied in the next section.

### Theorem 1

Let  $V = [N \ M] [\bar{P}^T \bar{P} - \bar{Q} \bar{Q}^T]^{-1} [N \ M]^T$  and  $W = P^{-1} M (P^{-1} M)^T$ .

1.  $\text{plim} (\hat{\sigma}^2 - \tilde{\sigma}^2) = 0$ , where  $\hat{\sigma}^2 = \epsilon^T V^{-1} \epsilon / T$  and  $\tilde{\sigma}^2 = \epsilon^T W^{-1} \epsilon / T$ .
2. If  $\lim_{T \rightarrow \infty} (X^T W^{-1} X / T)^{-1}$  exists, then  $\text{plim} (\hat{\beta} - \tilde{\beta}) = 0$ , with  $\hat{\beta}$  the Aitken estimator based on  $V$  and  $\tilde{\beta}$  the estimator based on  $W$ .

### Proof

See appendix.  $\square$

For the determinant of the covariance matrix we state the result in theorem 2. Asymptotically the value of the determinant of  $V$  tends not to that of  $W$ , which is equal to one, but it approaches the value of the determinant of an easy to compute ( $p \times p$ ) matrix.

### Theorem 2

If the invertibility condition for the MA part holds,

$$\lim_{T \rightarrow \infty} |V| = |I_p + (\underline{P}^T \underline{P} - \underline{Q} \underline{Q}^T)^{-1} (\underline{P} \underline{N} - \underline{M} \underline{Q})^T (\underline{M}^T \underline{M} - \underline{N} \underline{N}^T)^{-1} (\underline{P} \underline{N} - \underline{M} \underline{Q})|.$$

### Proof

Because of the structure of  $M$  and  $P$  we have  $|M| = |P| = 1$ . The determinant of the  $T \times T$  matrix  $V$  is  $|I_T + (PM^{-1}N - Q)(\underline{P}^T \underline{P} - \underline{Q} \underline{Q}^T)^{-1} (PM^{-1}N - Q)^T|$ , which is equal to the determinant of the  $p \times p$  matrix  $|I_p + (\underline{P}^T \underline{P} - \underline{Q} \underline{Q}^T)^{-1} (PM^{-1}N - Q)^T (PM^{-1}N - Q)|$ , as the second part of both expressions have the same non zero eigenvalues. Now,  $(PM^{-1}N - Q)^T (PM^{-1}N - Q) = (PN - MQ)^T (M^{-1})^T M^{-1} (PN - MQ)$ .

Here  $\underline{PN-MQ}$  is equal to  $\begin{bmatrix} \underline{PN-MQ} \\ 0 \end{bmatrix}$  because  $\underline{Q}$  and  $\underline{N}$  commute. This implies that  $\underline{M}^{-1}(\underline{PN-MQ})$  is equal to the first  $p$  columns of  $\underline{M}^{-1}$  times  $\underline{PN-MQ}$ . The elements of this column are  $(-\underline{M}^{-1}\underline{N})^{i-1}\underline{M}^{-1}$ ,  $i=1,2,\dots$ . What matters is thus only the upper left block of  $(\underline{M}^{-1})^T \underline{M}^{-1}$ , which is equal to

$\sum_{i=1} (\underline{M}^{-1})^T ((\underline{M}^{-1}\underline{N})^T)^{i-1} (\underline{M}^{-1}\underline{N})^{i-1} (\underline{M}^{-1})$ . Let this sum be  $\underline{S}$ . Then

$$\begin{aligned} \underline{M}^T \underline{S} \underline{M} &= \sum_{i=1} ((\underline{M}^{-1}\underline{N})^T)^{i-1} (\underline{M}^{-1}\underline{N})^{i-1} \\ &= \underline{I}_p + \sum_{i=2} \underline{N}^T (\underline{M}^{-1})^T ((\underline{M}^{-1}\underline{N})^T)^{i-2} (\underline{M}^{-1}\underline{N})^{i-2} \underline{M}^{-1} \underline{N} \\ &= \underline{I}_p + \underline{N}^T \left\{ \sum_{i=1} (\underline{M}^{-1})^T ((\underline{M}^{-1}\underline{N})^T)^{i-1} (\underline{M}^{-1}\underline{N})^{i-1} \underline{M}^{-1} \right\} \underline{N} \\ &= \underline{I}_p + \underline{N}^T \underline{S} \underline{N}. \end{aligned}$$

This equation has a solution if  $\underline{M}^T \otimes \underline{M}^T - \underline{N}^T \otimes \underline{N}^T$  is not singular. Its determinant, say  $\underline{D}$ , is

$$\begin{aligned} \underline{D} &= |\underline{M}^T \otimes \underline{M}^T - \underline{N}^T \otimes \underline{N}^T| \\ &= |\underline{I}_{p^2} - \underline{N}^T \otimes \underline{N}^T (\underline{M}^T \otimes \underline{M}^T)^{-1}| |\underline{M}^T \otimes \underline{M}^T| \\ &= |\underline{I}_{p^2} - (\underline{NM}^{-1})^T \otimes (\underline{NM}^{-1})^T|. \end{aligned}$$

The eigenvalues of  $\underline{NM}^{-1}$  are smaller than 1 in absolute value, implying that the determinant is non singular. Its solution  $\underline{S}$  is equal to  $(\underline{M}^T \underline{M} - \underline{NN}^T)^{-1}$  or  $(\underline{MM}^T - \underline{N}^T \underline{N})^{-1}$ . These expressions are equivalent as  $\underline{M}^T \underline{M} + \underline{N}^T \underline{N} = \underline{MM}^T + \underline{NN}^T$ : an equality which is shown easily by comparing the products of the commuting

matrices  $\begin{bmatrix} \underline{N} & \underline{M} \\ 0 & \underline{N} \end{bmatrix}^T$  and  $\begin{bmatrix} \underline{M} & 0 \\ \underline{N} & \underline{M} \end{bmatrix}$ . Then

$$\begin{aligned} \underline{S} &= (\underline{MM}^T - \underline{N}^T \underline{N})^{-1} \Rightarrow \underline{M}^T \underline{S} \underline{M} = (\underline{I}_p - \underline{M}^{-1} \underline{N}^T \underline{N} (\underline{M}^{-1})^T)^{-1}, \\ \underline{S} &= (\underline{M}^T \underline{M} - \underline{NN}^T)^{-1} \Rightarrow \underline{N}^T \underline{S} \underline{N} = (\underline{N}^{-1} \underline{M}^T \underline{M} (\underline{N}^{-1})^T - \underline{I}_p)^{-1} = ((\underline{M}^{-1} \underline{N}^T \underline{N} (\underline{M}^{-1})^T)^{-1} - \underline{I}_p)^{-1}, \text{ where the} \\ &\text{(commuting matrices) } \underline{M}^{-1} \text{ and } \underline{N}^T \text{ are interchanged. Direct verification} \\ &\text{concludes the proof. } \square \end{aligned}$$

For the pure AR-case we have  $\underline{M} = \underline{I}_p$  and  $\underline{N} = 0$ . The determinant becomes

$$|\underline{I}_p + (\underline{P}^T \underline{P} - \underline{QQ}^T)^{-1} \underline{QQ}^T| = |(\underline{P}^T \underline{P} - \underline{QQ}^T)^{-1} [\underline{P}^T \underline{P} - \underline{QQ}^T + \underline{Q}^T \underline{Q}]| = |(\underline{P}^T \underline{P} - \underline{QQ}^T)^{-1} \underline{PP}^T| =$$

$|(P^T P - Q Q^T)^{-1}|$ . For the pure MA-case we get a similar result:

$$|I_p + N^T (M^T M - N N^T)^{-1} N| = |M^T (M^T M - N N^T)^{-1} M| = |(M^T M - N N^T)^{-1}|.$$

## 5. Conclusion

Both inverse and determinant of the exact covariance matrix of ARMA distributed errors are rather complicated expressions. However, if the number of observations is large enough to permit an asymptotic approach we can use a simple form for the covariance matrix and its inverse. The value of the determinant used in the concentrated likelihood function reduces to the value of the determinant of an easy to compute matrix of order  $\max(p, q)$ .

## Appendix

### Proof of Theorem 1

1.  $\text{plim} (\hat{\sigma}^2 - \tilde{\sigma}^2) = 0$ , where  $\hat{\sigma}^2 = \varepsilon^T V^{-1} \varepsilon / T$  and  $\tilde{\sigma}^2 = \varepsilon^T W^{-1} \varepsilon / T$ .

Straightforward algebra gives:

$$\begin{aligned} E(\varepsilon^T W^{-1} \varepsilon - \varepsilon^T V^{-1} \varepsilon) &= E \text{tr}(\varepsilon^T W^{-1} \varepsilon - \varepsilon^T V^{-1} \varepsilon) \\ &= \text{tr} E(W^{-1} \varepsilon \varepsilon^T - V^{-1} \varepsilon \varepsilon^T) \\ &= \sigma^2 \text{tr}(W^{-1} V - I_T) \\ &= \sigma^2 \text{tr}(W^{-1} (W + (N - M P^{-1} Q) (P^T P - Q Q^T)^{-1} (N - M P^{-1} Q)^T) - I_T) \\ &= \sigma^2 \text{tr}(P^{-1} M (P^{-1} M)^T)^{-1} (N - M P^{-1} Q) (P^T P - Q Q^T)^{-1} (N - M P^{-1} Q)^T \\ &= \sigma^2 \text{tr}((P^{-1} M)^T)^{-1} (P^{-1} M)^{-1} (N - M P^{-1} Q) (P^T P - Q Q^T)^{-1} (N - M P^{-1} Q)^T \\ &= \sigma^2 \text{tr}(P^{-1} M)^{-1} (N - M P^{-1} Q) (P^T P - Q Q^T)^{-1} (N - M P^{-1} Q)^T ((P^{-1} M)^T)^{-1} \\ &= \sigma^2 \text{tr}(P M^{-1} N - Q) (P^T P - Q Q^T)^{-1} (P M^{-1} N - Q)^T. \end{aligned}$$

In view of lemma 2 this expression divided by  $T$  vanishes for  $T \rightarrow \infty$ .  $\square$

2. If  $\lim_{T \rightarrow \infty} (X^T W^{-1} X / T)^{-1}$  exists, then  $\text{plim} (\hat{\beta} - \tilde{\beta}) = 0$ , with  $\hat{\beta}$  the Aitken

estimator based on  $V$  and  $\tilde{\beta}$  the Aitken estimator based on  $W$ .

This part needs more manipulation. Write the difference between  $\hat{\beta}$  and  $\tilde{\beta}$  as a function of the error  $\epsilon$ :

$$\hat{\beta} = (X^T V^{-1} X)^{-1} X V^{-1} y = (X^T V^{-1} X)^{-1} X V^{-1} (X \beta + \epsilon) = \beta + (X^T V^{-1} X)^{-1} X V^{-1} \epsilon \text{ and}$$

$$\tilde{\beta} = \beta + (X^T W^{-1} X)^{-1} X W^{-1} \epsilon, \text{ which gives for the difference}$$

$$\hat{\beta} - \tilde{\beta} = ((X^T V^{-1} X)^{-1} X V^{-1} - (X^T W^{-1} X)^{-1} X W^{-1}) \epsilon \text{ or } Z \epsilon.$$

Observe that  $E \|Z \epsilon\|^2 = \text{tr } E Z \epsilon \epsilon^T Z^T = \sigma^2 \text{tr } Z V Z^T$ . We will show, that  $\lim_{T \rightarrow \infty} E \|Z \epsilon\|^2 = 0$ ,

which implies  $\text{plim}(Z \epsilon) = \text{plim}(\hat{\beta} - \tilde{\beta}) = 0$ , by proving that  $\text{tr } Z^T V Z$  has an upper limit similar to the expression in lemma 2. We rewrite  $\hat{\beta} - \tilde{\beta}$  replacing  $V$  by  $W$  and several other symbols. Introduce

$$A = N - P^{-1} M Q, \quad (3)$$

$$\Delta = P^T P - Q Q^T, \quad (4)$$

$$H = A^T W^{-1} A + \Delta \quad (5)$$

and

$$G = A^T W^{-1} X (X^T W^{-1} X)^{-1} X^T W^{-1} A - H. \quad (6)$$

Then  $V = W + A \Delta^{-1} A^T$ , using (3) and (4). This gives for the inverse, using a well known formula for the inverse of the sum of two matrices and (5),

$$\begin{aligned} V^{-1} &= (W + A \Delta^{-1} A^T)^{-1} \\ &= W^{-1} - W^{-1} A (A^T W^{-1} A + \Delta)^{-1} A^T W^{-1} \end{aligned}$$

or

$$V^{-1} = W^{-1} - W^{-1} A H^{-1} A^T W^{-1}. \quad (7)$$

Next express  $(X^T V^{-1} X)^{-1}$  in  $W$  and  $A$ :

$$\begin{aligned} (X^T V^{-1} X)^{-1} &= (X^T W^{-1} X - X^T W^{-1} A H^{-1} A^T W^{-1} X)^{-1} \\ &= (X^T W^{-1} X)^{-1} - (X^T W^{-1} X)^{-1} X^T W^{-1} A (A^T W^{-1} X (X^T W^{-1} X)^{-1} X^T W^{-1} A - H)^{-1} A^T W^{-1} X (X^T W^{-1} X)^{-1} \end{aligned}$$

or using (6),

$$(X^T V^{-1} X)^{-1} = (X^T W^{-1} X)^{-1} - (X^T W^{-1} X)^{-1} X^T W^{-1} A G^{-1} A^T W^{-1} X (X^T W^{-1} X)^{-1}. \quad (8)$$



Note that G is negative:

$$\begin{aligned}
 G &= A^T W^{-1} X (X^T W^{-1} X)^{-1} X^T W^{-1} A - H \\
 &= A^T W^{-1} X (X^T W^{-1} X)^{-1} X^T W^{-1} A - A^T W^{-1} A - \Delta \\
 &= A^T (W^{-1} X (X^T W^{-1} X)^{-1} X^T W^{-1} A - W^{-1}) A - \Delta \\
 &= -A^T (W^{-1} - W^{-1} X (X^T W^{-1} X)^{-1} X^T W^{-1}) A - \Delta.
 \end{aligned}$$

Furthermore we need the expression  $W^{-1} V W^{-1}$  rewritten in A and W:

$$W^{-1} V W^{-1} = W^{-1} (W + A \Delta^{-1} A^T) W^{-1}$$

or

$$W^{-1} V W^{-1} = W^{-1} + W^{-1} A \Delta^{-1} A^T W^{-1}. \quad (9)$$

Finally express  $Z V Z^T$  in X and W:

$$\begin{aligned}
 Z V Z^T &= ((X^T V^{-1} X)^{-1} X V^{-1} - (X^T W^{-1} X)^{-1} X W^{-1}) V ((X^T V^{-1} X)^{-1} X V^{-1} - (X^T W^{-1} X)^{-1} X W^{-1})^T \\
 &= (X^T V^{-1} X)^{-1} X V^{-1} V V^{-1} X^T (X^T V^{-1} X)^{-1} - (X^T V^{-1} X)^{-1} X V^{-1} V W^{-1} X^T (X^T W^{-1} X)^{-1} - \\
 &\quad (X^T W^{-1} X)^{-1} X W^{-1} V V^{-1} X^T (X^T V^{-1} X)^{-1} + (X^T W^{-1} X)^{-1} X W^{-1} V W^{-1} X^T (X^T W^{-1} X)^{-1} \\
 &= -(X^T V^{-1} X)^{-1} + (X^T W^{-1} X)^{-1} X W^{-1} V W^{-1} X^T (X^T W^{-1} X)^{-1}.
 \end{aligned}$$

Use (8) and (9) to replace  $(X^T V^{-1} X)^{-1}$  and  $W^{-1} V W^{-1}$ :

$$\begin{aligned}
 Z V Z^T &= -(X^T W^{-1} X)^{-1} + (X^T W^{-1} X)^{-1} X^T W^{-1} A G^{-1} A^T W^{-1} X (X^T W^{-1} X)^{-1} + \\
 &\quad (X^T W^{-1} X)^{-1} X (W^{-1} + W^{-1} A \Delta^{-1} A^T W^{-1}) X^T (X^T W^{-1} X)^{-1} \\
 &= -(X^T W^{-1} X)^{-1} + (X^T W^{-1} X)^{-1} X^T W^{-1} A G^{-1} A^T W^{-1} X (X^T W^{-1} X)^{-1} + \\
 &\quad (X^T W^{-1} X)^{-1} + (X^T W^{-1} X)^{-1} X W^{-1} A \Delta^{-1} A^T W^{-1} X^T (X^T W^{-1} X)^{-1} \\
 &= (X^T W^{-1} X)^{-1} X^T W^{-1} A (G^{-1} + \Delta^{-1}) A^T W^{-1} X (X^T W^{-1} X)^{-1} \\
 &< (X^T W^{-1} X)^{-1} X^T W^{-1} A \Delta^{-1} A^T W^{-1} X (X^T W^{-1} X)^{-1},
 \end{aligned}$$

because G is definite negative.

Since we assume that the limit of the inverse of

$$X^T W^{-1} X / T = X^T (P^{-1} M (P^{-1} M)^T)^{-1} X / T = (M^{-1} P X / \sqrt{T})^T (M^{-1} P X / \sqrt{T}) \text{ exists, } \lim_{T \rightarrow \infty} (M^{-1} P X / \sqrt{T})$$

$$\text{has to exist. Write } \Xi \text{ for } \left[ \frac{X^T W^{-1} X}{T} \right]^{-1} \left[ \frac{(M^{-1} P X)^T}{\sqrt{T}} \right].$$



$$\text{As } \frac{X^T W^{-1} A}{T} = \frac{(M^{-1} P X)^T M^{-1} P (N - P^{-1} M Q)}{T} = \left( \frac{(M^{-1} P X)^T}{\sqrt{T}} \right) \left( \frac{P M^{-1} N - Q}{\sqrt{T}} \right) \text{ we get}$$

$$\begin{aligned} \lim_{T \rightarrow \infty} \text{tr} Z V Z^T &< \text{tr} \lim_{T \rightarrow \infty} \left( \frac{X^T W^{-1} X}{T} \right)^{-1} \left( \frac{X^T W^{-1} A}{T} \right) \Delta^{-1} \left( \frac{A^T W^{-1} X}{T} \right) \left( \frac{X^T W^{-1} X}{T} \right)^{-1} \\ &< \text{tr} \lim_{T \rightarrow \infty} \Xi \{ (P M^{-1} N - Q) / \sqrt{T} \} \Delta^{-1} \{ (P M^{-1} N - Q)^T / \sqrt{T} \} \Xi^T \\ &< \text{tr} \lim_{T \rightarrow \infty} \Xi \lim_{T \rightarrow \infty} \{ (P M^{-1} N - Q) \Delta^{-1} (P M^{-1} N - Q)^T / T \} \lim_{T \rightarrow \infty} \Xi^T \\ &= 0, \text{ in view of lemma 2. } \square \end{aligned}$$

## References

- Judge, G.C. et al. (1985), *The Theory and Practice of Econometrics*, New York.
- Van der Leeuw, J.L. (1992), The Covariance Matrix of ARMA-errors in Closed Form, *Research Memorandum Tilburg University*, 562.
- Van der Leeuw, J.L. (1993), First Order Conditions for the Maximum Likelihood Estimation of an Exact ARMA Model, *Research Memorandum Tilburg University*, 611.
- Pagan, A (1974) A Generalised Approach to the Treatment of Autocorrelation. *Australian Economic Papers*, 13.
- Magnus, J.R. (1978), Maximum Likelihood Estimation of the GLS Model with Unknown Parameters in the Disturbance Covariance Matrix, *Journal of Econometrics*, 7.
- Zinde-Walsh, V. and J.W. Galbraith (1991), Estimation of a Linear Regression Model with Stationary ARMA(p,q) Errors, *Journal of Econometrics*, 47.

## IN 1993 REEDS VERSCHENEN

- 588 Rob de Groof and Martin van Tuijl  
The Twin-Debt Problem in an Interdependent World  
Communicated by Prof.dr. Th. van de Klundert
- 589 Harry H. Tigelaar  
A useful fourth moment matrix of a random vector  
Communicated by Prof.dr. B.B. van der Genugten
- 590 Niels G. Noorderhaven  
Trust and transactions; transaction cost analysis with a differential behavioral assumption  
Communicated by Prof.dr. S.W. Douma
- 591 Henk Roest and Kitty Koelemeijer  
Framing perceived service quality and related constructs A multilevel approach  
Communicated by Prof.dr. Th.M.M. Verhallen
- 592 Jacob C. Engwerda  
The Square Indefinite LQ-Problem: Existence of a Unique Solution  
Communicated by Prof.dr. J. Schumacher
- 593 Jacob C. Engwerda  
Output Deadbeat Control of Discrete-Time Multivariable Systems  
Communicated by Prof.dr. J. Schumacher
- 594 Chris Veld and Adri Verboven  
An Empirical Analysis of Warrant Prices versus Long Term Call Option Prices  
Communicated by Prof.dr. P.W. Moerland
- 595 A.A. Jeunink en M.R. Kabir  
De relatie tussen aandeelhoudersstructuur en beschermingsconstructies  
Communicated by Prof.dr. P.W. Moerland
- 596 M.J. Coster and W.H. Haemers  
Quasi-symmetric designs related to the triangular graph  
Communicated by Prof.dr. M.H.C. Paardekooper
- 597 Noud Gruijters  
De liberalisering van het internationale kapitaalverkeer in historisch-institutioneel perspectief  
Communicated by Dr. H.G. van Gemert
- 598 John Görtzen en Remco Zwetheul  
Weekend-effect en dag-van-de-week-effect op de Amsterdamse effectenbeurs?  
Communicated by Prof.dr. P.W. Moerland
- 599 Philip Hans Franses and H. Peter Boswijk  
Temporal aggregation in a periodically integrated autoregressive process  
Communicated by Prof.dr. Th.E. Nijman

- 600 René Peeters  
On the p-ranks of Latin Square Graphs  
Communicated by Prof.dr. M.H.C. Paardekooper
- 601 Peter E.M. Borm, Ricardo Cao, Ignacio García-Jurado  
Maximum Likelihood Equilibria of Random Games  
Communicated by Prof.dr. B.B. van der Genugten
- 602 Prof.dr. Robert Bannink  
Size and timing of profits for insurance companies. Cost assignment for products with multiple deliveries.  
Communicated by Prof.dr. W. van Hulst
- 603 M.J. Coster  
An Algorithm on Addition Chains with Restricted Memory  
Communicated by Prof.dr. M.H.C. Paardekooper
- 604 Ton Geerts  
Coordinate-free interpretations of the optimal costs for LQ-problems subject to implicit systems  
Communicated by Prof.dr. J.M. Schumacher
- 605 B.B. van der Genugten  
Beat the Dealer in Holland Casino's Black Jack  
Communicated by Dr. P.E.M. Borm
- 606 Gert Nieuwenhuis  
Uniform Limit Theorems for Marked Point Processes  
Communicated by Dr. M.R. Jaïbi
- 607 Dr. G.P.L. van Roij  
Effectisering op internationale financiële markten en enkele gevolgen voor banken  
Communicated by Prof.dr. J. Sijben
- 608 R.A.M.G. Joosten, A.J.J. Talman  
A simplicial variable dimension restart algorithm to find economic equilibria on the unit simplex using  $n(n+1)$  rays  
Communicated by Prof.Dr. P.H.M. Ruys
- 609 Dr. A.J.W. van de Gevel  
The Elimination of Technical Barriers to Trade in the European Community  
Communicated by Prof.dr. H. Huizinga
- 610 Dr. A.J.W. van de Gevel  
Effective Protection: a Survey  
Communicated by Prof.dr. H. Huizinga
- 611 Jan van der Leeuw  
First order conditions for the maximum likelihood estimation of an exact ARMA model  
Communicated by Prof.dr. B.B. van der Genugten

- 612 Tom P. Faith  
 Bertrand-Edgeworth Competition with Sequential Capacity Choice  
 Communicated by Prof.Dr. S.W. Douma
- 613 Ton Geerts  
 The algebraic Riccati equation and singular optimal control: The discrete-time case  
 Communicated by Prof.dr. J.M. Schumacher
- 614 Ton Geerts  
 Output consistency and weak output consistency for continuous-time implicit systems  
 Communicated by Prof.dr. J.M. Schumacher
- 615 Stef Tijs, Gert-Jan Otten  
 Compromise Values in Cooperative Game Theory  
 Communicated by Dr. P.E.M. Borm
- 616 Dr. Pieter J.F.G. Meulendijks and Prof.Dr. Dick B.J. Schouten  
 Exchange Rates and the European Business Cycle: an application of a 'quasi-empirical' two-country model  
 Communicated by Prof.Dr. A.H.J.J. Kolnaar
- 617 Niels G. Noorderhaven  
 The argumentational texture of transaction cost economics  
 Communicated by Prof.Dr. S.W. Douma
- 618 Dr. M.R. Jaïbi  
 Frequent Sampling in Discrete Choice  
 Communicated by Dr. M.H. ten Raai
- 619 Dr. M.R. Jaïbi  
 A Qualification of the Dependence in the Generalized Extreme Value Choice Model  
 Communicated by Dr. M.H. ten Raai
- 620 J.J.A. Moors, V.M.J. Coenen, R.M.J. Heuts  
 Limiting distributions of moment- and quantile-based measures for skewness and kurtosis  
 Communicated by Prof.Dr. B.B. van der Genugten
- 621 Job de Haan, Jos Benders, David Bennett  
 Symbiotic approaches to work and technology  
 Communicated by Prof.dr. S.W. Douma
- 622 René Peeters  
 Orthogonal representations over finite fields and the chromatic number of graphs  
 Communicated by Dr.ir. W.H. Haemers
- 623 W.H. Haemers, E. Spence  
 Graphs Cospectral with Distance-Regular Graphs  
 Communicated by Prof.dr. M.H.C. Paardekooper

- 624 Bas van Aarle  
The target zone model and its applicability to the recent EMS crisis  
Communicated by Prof.dr. H. Huizinga
- 625 René Peeters  
Strongly regular graphs that are locally a disjoint union of hexagons  
Communicated by Dr.ir. W.H. Haemers
- 626 René Peeters  
Uniqueness of strongly regular graphs having minimal  $p$ -rank  
Communicated by Dr.ir. W.H. Haemers
- 627 Freek Aertsen, Jos Benders  
Tricks and Trucks: Ten years of organizational renewal at DAF?  
Communicated by Prof.dr. S.W. Douma
- 628 Jan de Klein, Jacques Roemen  
Optimal Delivery Strategies for Heterogeneous Groups of Porkers  
Communicated by Prof.dr. F.A. van der Duyn Schouten
- 629 Imma Curiel, Herbert Hamers, Jos Potters, Stef Tijs  
The equal gain splitting rule for sequencing situations and the general nucleolus  
Communicated by Dr. P.E.M. Borm
- 630 A.L. Hempenius  
Een statische theorie van de keuze van bankrekening  
Communicated by Prof.Dr.Ir. A. Kapteyn
- 631 Cok Vrooman, Piet van Wijngaarden, Frans van den Heuvel  
Prevention in Social Security: Theory and Policy Consequences  
Communicated by Prof.Dr. A. Kolnaar



## IN 1994 REEDS VERSCHENEN

- 632 B.B. van der Genugten  
Identification, estimating and testing in the restricted linear model  
Communicated by Dr. A.H.O. van Soest
- 633 George W.J. Hendrikse  
Screening, Competition and (De)Centralization  
Communicated by Prof.dr. S.W. Douma
- 634 A.J.T.M. Weeren, J.M. Schumacher, and J.C. Engwerda  
Asymptotic Analysis of Nash Equilibria in Nonzero-sum Linear-Quadratic Differential Games. The Two-Player case  
Communicated by Prof.dr. S.H. Tijs
- 635 M.J. Coster  
Quadratic forms in Design Theory  
Communicated by Dr.ir. W.H. Haemers
- 636 Drs. Erwin van der Krabben, Prof.dr. Jan G. Lambooy  
An institutional economic approach to land and property markets - urban dynamics and institutional change  
Communicated by Dr. F.W.M. Boekema
- 637 Bas van Aarle  
Currency substitution and currency controls: the Polish experience of 1990  
Communicated by Prof.dr. H. Huizinga
- 638 J. Bell  
Joint Ventures en Ondernemerschap: Interpreneurship  
Communicated by Prof.dr. S.W. Douma
- 639 Frans de Roon and Chris Veld  
Put-call parities and the value of early exercise for put options on a performance index  
Communicated by Prof.dr. Th.E. Nijman
- 640 Willem J.H. Van Groenendaal  
Assessing demand when introducing a new fuel: natural gas on Java  
Communicated by Prof.dr. J.P.C. Kleijnen
- 641 Henk van Gemert & Noud Gruijters  
Patterns of Financial Change in the OECD area  
Communicated by Prof.dr. J.J. Sijben
- 642 Drs. M.R.R. van Bremen, Drs. T.A. Marra en Drs. A.H.F. Verboven  
Aardappelen, varkens en de termijnhandel: de reële optietheorie toegepast  
Communicated by Prof.dr. P.W. Moerland



- 643 W.J.H. Van Groenendaal en F. De Gram  
The generalization of netback value calculations for the determination of industrial demand for natural gas  
Communicated by Prof.dr. J.P.C. Kleijnen
- 644 Karen Aardal, Yves Pochet and Laurence A. Wolsey  
Capacitated Facility Location: Valid Inequalities and Facets  
Communicated by Dr.ir. W.H. Haemers
- 645 Jan J.G. Lemmen  
An Introduction to the Diamond-Dybvig Model (1983)  
Communicated by Dr. S. Eijffinger
- 646 Hans J. Gremmen and Eva van Deurzen-Mankova  
Reconsidering the Future of Eastern Europe: The Case of Czecho-Slovakia  
Communicated by Prof.dr. H.P. Huizinga
- 647 H.M. Webers  
Non-uniformities in spatial location models  
Communicated by Prof.dr. A.J.J. Talman
- 648 Bas van Aarle  
Social welfare effects of a common currency  
Communicated by Prof.dr. H. Huizinga
- 649 Laurence A.G.M. van Lent  
De winst is absoluut belangrijk!  
Communicated by Prof.drs. G.G.M. Bak
- 650 Bert Hamminga  
Jager over de theorie van de internationale handel  
Communicated by Prof.dr. H. Huizinga
- 651 J.Ch. Caanen and E.N. Kertzman  
A comparison of two methods of inflation adjustment  
Communicated by Prof.dr. J.A.G. van der Geld
- 652 René van den Brink  
A Note on the  $\tau$ -Value and  $\tau$ -Related Solution Concepts  
Communicated by Prof.dr. P.H.M. Ruys
- 653 J. Engwerda and G. van Willigenburg  
Optimal sampling-rates of digital LQ and LQG tracking controllers with costs associated to sampling  
Communicated by Prof.dr. J.M. Schumacher
- 654 J.C. de Vos  
A Thousand Golden Ten Orbits  
Communicated by Prof.dr. B.B. van der Genugten

- 655 Gert-Jan Otten, Peter Borm, Stef Tijs  
A Note on the Characterizations of the Compromise Value  
Communicated by Prof.dr. A.J.J. Talman
- 656 René Peeters  
On the p-ranks of the adjacency matrices of distance-regular graphs  
Communicated by Dr.ir. W.H. Haemers
- 657 J. Kriens, L.W.G. Strijbosch, J. Vörös  
Differentiability Properties of the Efficient  $(\mu, \sigma^2)$ -Set in the Markowitz Portfolio Selection Method  
Communicated by Prof.dr. A.J.J. Talman
- 658 Prof.dr. E.J. Bijnen and dr. M.F.C.M. Wijn  
Corporate Prediction Models, Ratios or Regression Analysis?  
Communicated by Prof.dr. W. van Hulst
- 659 Edwin R. van Dam  
Regular graphs with four eigenvalues  
Communicated by Dr.ir. W.H. Haemers
- 660 G.J. van der Pijl  
Quality of information and the goals and targets of the organization:  
a model and a method  
Communicated by Prof.dr. P.M.A. Ribbers
- 661 Jan van der Leeuw  
The second derivative of the likelihood of an exact ARMA model  
Communicated by Dr. H.H. Tigelaar

Bibliotheek K. U. Brabant



17 000 01126800 1

Katholieke Universiteit Brabant  
PO Box 90153  
5000 LE Tilburg  
The Netherlands